



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A01N 45/00, 25/30 // (A01N 45/00, 57:20, 45:00, 43:40, 37:42, 33:12)	A1	(11) International Publication Number: WO 00/02454 (43) International Publication Date: 20 January 2000 (20.01.00)
(21) International Application Number: PCT/EP99/04592 (22) International Filing Date: 2 July 1999 (02.07.99) (30) Priority Data: 98112541.2 7 July 1998 (07.07.98) EP (71) Applicant (for all designated States except US): BASF AKTIENGESELLSCHAFT [DE/DE]; D-67056 Ludwigshafen (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): RADEMACHER, Wilhelm [DE/DE]; Austrasse 1, D-67117 Limburgerhof (DE). SCHNEIDER, Karl-Heinrich [DE/DE]; Am Kelleracker 20, D-67271 Kleinkarlbach (DE). KOBER, Reiner [DE/DE]; Im Schlittweg 20, D-67136 Fußgönheim (DE). KING, Roderick, Whitfield [AU/AU]; 22 Norman Street, Deakin, ACT 2600 (AU). MANDER, Lewis, Norman [AU/AU]; 15 Juwin Street, Aranda, ACT 2614 (AU). PHARIS, Richard, Persons [CA/CA]; RR2 Box 609, Cochrane, Alberta (CA). (74) Common Representative: BASF AKTIENGESELLSCHAFT; D-67056 Ludwigshafen (DE).		(81) Designated States: AL, AU, BG, BR, BY, CA, CN, CZ, GE, HR, HU, ID, IL, IN, JP, KR, KZ, LT, LV, MK, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, ZA, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PLANT GROWTH REGULATING FORMULATIONS (57) Abstract Plant growth regulation preparations comprising a) 0.1 to 20 % by weight of a 16,17-dihydro gibberrelin of general formulae (Ia) or (Ib); b) up to 99.9 % by weight of a formulation additive selected from the group consisting of b1) the reaction products of triglycerides based on carboxylic acids having 2 to 30 carbon atoms and ethylene oxide and/or propylene oxide in the presence of a base, and/or b2) fatty acid esters of sugar alcohol polyethoxylates; c) up to 50 % by weight of an organic solvent; d) 0.1 to 50 % by weight of a formulation auxiliary different from b1 and b2; e) up to 50 % by weight of additional plant growth regulating compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Plant growth regulating formulations

The present invention relates to plant growth regulating preparations comprising

5

a) 0.1 to 20 % by weight of a 16,17-dihydro gibberrelin of general formulae Ia or Ib

10 b) up to 99.9 % by weight of an formulation additive selected from the group consisting of

b1) the reaction products of triglycerides based on carboxylic acids having 2 to 30 carbon atoms and ethylene oxide and/or propylene oxide in the presence of a base, and/or
15

b2) fatty acid esters of sugar alcohol polyethoxylates

20 c) up to 50 % by weight of an organic solvent

d) 0.1 to 50 % by weight of a formulation auxiliary different from b1 and b2,

25 e) up to 50 % by weight of additional plant growth regulating compounds.

Furthermore, the present invention relates to the use of the said preparations for the regulation of plant growth.

30 Numerous plant growth substances are known. Part of these substances are used in agriculture and horticulture in order to induce desired effects, for instance on seed germination and seedling growth

- rooting
- 35 • dormancy
- juvenility, maturity and senescence
- flowering
- abscission of leaves, flowers and fruits
- fruit set and development
- 40 • tuber formation
- growth of shoot and root
- photoassimilation
- control of unwanted plants
- senescence of whole plants or single organs

45

[cf. R. N. Arteca (1996) Plant Growth Substances, Chapman & Hall, New York.]

2

- Gibberellins (GAs) represent one group of plant growth substances. GAs occur naturally in higher plants where they function as phytohormones being involved in the regulation of growth and flowering and numerous other developmental processes. Certain GAs, particularly gibberellic acid or GA₃ (ent-3 α ,10,13-trihydroxy-20-norgiberrell-1,16-diene-7,19-dioic acid 19,10-lactone), are commercially available as fermentation products of the fungus *Gibberella fujikuroi*.
- 10 C-16,17-dihydro GAs and related structures have been known since many years. They either occur naturally in higher plants (e.g. GA₈₂ and GA₈₃) or in the fungi *Gibberella fujikuroi* (e.g. GA₂, GA₁₀, GA₄₂) and *Phaeosphaeria sp.* (e.g. GA₈₂).
- 15 WO 93/03616 and PCT/AU95/000528 describe for the first time the biological activity of known and newly synthesized 16,17-dihydro GAs and their potential utility for agricultural, horticultural and plant breeding applications. In particular the following biological effects have been described as features of the
- 20 16,17-dihydro GAs and/or other ring D-modified GAs:
- dwarfing
 - growth retardation of stem and shoot and/or root
 - promotion of bud break
 - flowering
 - 25 • improved fruit quality
 - inhibition of fruit ripening
 - prevention of whole plant and/or organ senescence
 - improved fruit set
 - control of weed growth
 - 30 • inducing male sterility
 - retarded bud break
 - tillering

- Further details on chemical and biological aspects of distinct
- 35 16,17-dihydro or other ring D-modified GAs have been published in the recent scientific literature.

- A major obstacle to introduce the ring D-modified gibberellin derivatives described in WO 93/03616 and PCT/AU95/000528 into
- 40 agricultural and horticultural practice is caused by the costs required to produce these compounds. The production costs are primarily affected by the price of the starting material GA₃ [cf. W. Rademacher (1997): Gibberellins, in Fungal Biotechnology, T. Anke (ed.), Chapman & Hall, Weinheim, pp. 193-205].
- 45

3

There thus exists a need in formulations based on ring modified D-gibberelin derivatives having improved biological activity and thus better economical feasibility.

5 The object of the present invention was to provide plant growth regulating preparations based on ring D-modified gibberellin derivatives having a very good biological activity, in particular at low application rates.

10 This object has been achieved with the plant growth regulating formulations as defined in claim 1 and hereinbefore.


Preferred embodiments of the instant invention are set forth in the dependent claims and the following detailed specification.

15

The C-16,17-dihydro gibberellins (component a) have the chemical formulae Ia or Ib

20

25




Ia

30 wherein A, B, C, D, E and F independently represent hydrogen atoms or hydroxyl groups and the dotted line represents one optional double bond either between the carbon atoms in position 1 and 2 or between the carbon atoms in positions 2 and 3, and

35

40



Ib

45 wherein

4

R¹ represents H or OH, OC(=O)R⁵ or OR⁵


R², R³, which may be the same or different, each represent H, F, Cl, Br, lower (C₁₋₆) alkyl, lower (C₂₋₆) alkenyl, lower (C₃₋₆) cycloalkyl, or CH₂X (wherein X is F, Cl or Br);

R⁴ indicates that the A ring may be (i) unfunctionalised, or (ii) contain a 1,2-double bond or 2,3-double bond, or (iii) contain a 3 α - or 3 β -OH, F, Cl or Br group with or without a 1,2-double bond, or (iv) contain a 1 α - or 1 β -OH, F Cl or Br group with or without a 2,3-double bond;

R⁵ represents C₁-C₆-alkyl.

The following compounds are particularly preferred in the preparation of the inventive preparations:

20




II

25

exo-16,17-dihydro GA₅

30




III

35

endo-16,17-Dihydro-GA₅


40




IV

45


16,17-Dichloromethano-dihydro-GA₅




10
16,17-Dichloromethano-
dihydro-GA₅-13-acetate



endo-16,17-Dihydro-GA₅-13-acetate




35
exo-16,17-Dihydro-GA₅-13-acetate



endo-16,17-Dihydro-GA₅-13-n-propyl ether

6

5



IX

exo-16,17-Dihydro-GA₅-13-n-propyl ether

10

The content of the component a) in the inventive formulations is in the range of from 0.1 to 20, in particular 0.2 to 10 % by weight of the formulation.

15

As component b) the inventive formulations contain

- b) up to 99.9 % by weight of an formulation additive selected from the group consisting of

20

- b1) the reaction products of triglycerides based on carboxylic acids having 2 to 30 carbon atoms and ethylene oxide and/or propylene oxide in the presence of a base, and/or

25

- b2) fatty acid esters of sugar alcohol polyethoxylates.

The component b1 is obtainable by the reaction of an oil/fat based on a triglyceride of carboxylic acids having 2 to 30 carbon atoms and ethylene oxide and/or propylene oxide in the presence of a base. Fatty acid alkoxyates are primarily formed.

30

In the triglycerides, three equivalents of carboxylic acid are esterified with glycerol. The carboxylic acids can be saturated or mono- or polyethylenically unsaturated.

35

Preferably, the present invention uses naturally occurring oils and fats which contain triglycerides as the main constituent. They can be crude, denatured or refined.

40

Suitable natural oils and fats are: vegetable oils such as olive oil, safflower oil, soybean oil, groundnut oil, cotton oil, corn oil, rape oil, castor oil, sunflower oil, coffee oil, linseed oil, coconut fat and mixtures thereof, animal fats and oils such as fish oils, e.g. sardine oil, herring oil, salmon oil, shark-liver oil or whale oil, and further tallow oil, bone oil, woofat fractions and bovine tallow.

45

The following vegetable oils are preferred: castor oil, soybean oil, rape oil and corn oil. Particularly preferred of these is castor oil.

- 5 Per mol of the triglyceride on which the oil is based, from 1 to 100, preferably from 10 to 35 and in particular from 15 to 30, mol of ethylene oxide and/or propylene oxide are employed.

Suitable bases are especially inorganic bases such as the alkali
10 metal or alkaline earth metal hydroxides, e.g. sodium hydroxide, potassium hydroxide or calcium hydroxide. Instead of the direct use of hydroxides, it is possible to use carbonates or hydro-talcites, which, if appropriate, were hydrophobized with
15 22 carbon atoms or the ethoxylates of alcohols of this type, or basic alkaline earth metal phosphates such as strontium phosphate, barium phosphate and calcium phosphate, each of which also form hydroxide ions in the presence of small amounts of water.

20

The base particularly used is potassium hydroxide.

- Per mol of triglyceride, from 0.1 to 5, and in particular from 0.1 to 2, % by weight of base, based on the weight of the
25 triglyceride, are generally added to the reaction mixture.

The reaction is generally carried out at elevated pressure, preferably at from 1 to 10 and in particular at from 1 to 5 bar and at from 80 to 230 and especially from 100 to 150°C.

30

Working-up is in general carried out such that excess ethylene oxide or propylene oxide is removed under reduced pressure.

- Otherwise, the carrying-out of such alkoxylation reactions and
35 the isolation of the resulting reaction products is known to the person skilled in the art (cf. N. Schönfeldt, Grenzflächenaktive Ethylenoxidaddukte [Surface-active ethylene oxide adducts], Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, 1984).

- 40 A particularly preferred product for component b1 is the product obtainable by the reaction of an oil/fat based on a triglyceride of carboxylic acids having 10 to 20 carbon atoms and 10 to 30 mol equivalents of ethylene oxide and/or propylene oxide in the presence of hydroxide ions.

45

8

Preferred as component b2 are the esters derived from C₈-C₂₈ fatty acids with C₅- and/or C₆ sugar alcohol polyethoxylates comprising preferably of from 10 to 100, particularly 20 to 50 ethoxylate units.

5

Preferred fatty acids are oleic acid, stearic acid, pelargonic acid and 2-Ethylhexanoic acid or mixtures thereof.

Component b can be solely constituted of component b1 or b2 or
10 of mixtures of both species.

Component b constitutes up to 99.9, preferably 0.5 to 50, in particular 0.5 to 25 % by weight of the formulations in accordance with the instant invention.

15

The formulations according to the invention can moreover additionally contain further customary additives such as surfactants, antifoams, cosolvents, etc.

20 Suitable surfactants are:

- anionic surfactants, e.g. alkali metal, alkaline earth metal or ammonium salts of the fatty acids such as potassium stearate, alkylsulfates, alkyl ether sulfates, alkyl- or
25 isoalkylsulfonates, alkylbenzenesulfonates such as Na dodecylbenzenesulfonates, alkyl naphthalenesulfonates, alkylmethyl ester sulfonates, acylglutamates, alkylsuccinic acid ester sulfonates, sarcosinates such as sodium lauroyl sarcosinate or taurates,
- 30 - cationic surfactants, e.g. alkyltrimethylammonium halides or alkylsulfates, alkylpyridinium halides or dialkyldimethylammonium halides or alkylsulfates,
- 35 - nonionic surfactants, e.g. alkoxylated animal or vegetable fats and oils such as corn oil ethoxylates, castor oil ethoxylates, tallow fat ethoxylates, glycerol esters such as glycerol monostearate, fatty alcohol alkoxylates and oxo-alcohol alkoxylates, fatty acid alkoxylates such as oleic
40 acid ethoxylates, alkylphenyl alkoxylates such as isononylphenol ethoxylates, fatty amine alkoxylates, fatty acid amide alkoxylates, sugar surfactants such as sorbitan fatty acid esters (sorbitan monooleate, sorbitan tristearate), polyoxyethylenesorbitan fatty acid esters, alkyl polyglycosides,
45 N-alkylgluconamides, alkylmethyl sulfoxides, alkyl dimethylphosphine oxides such as tetradecyldimethylphosphine oxide,

- zwitterionic surfactants, e.g. sulfobetaines, carboxybetaines, alkyldimethylamine oxides such as tetradecyldimethylamine oxide,
- 5 - polymer surfactants, e.g. di-, tri- or multi-blockpolymers of the type $(AB)_x$, ABA and BAB such as polyethylene oxide-block-polypropylene oxide, polystyrene-block-polyethylene oxide, AB comb polymers such as polymethacrylate or polyacrylate comb-polyethylene oxide,
- 10 - perfluoro surfactants, silicone surfactants, phospholipids such as lecithin, amino acid surfactants such as N-lauroyl glutamate, surface-active homo- and copolymers such as polyvinylpyrrolidone, polyacrylic acid, polyvinyl alcohol,
- 15 polyethylene oxide, maleic anhydride-isobutene copolymers, vinylpyrrolidone-vinyl acetate copolymers.

Preferably, the surfactant used is one or more homogeneous or mixed esters of phosphoric acid or diphosphoric acid with polyalkylene oxide ethers, the polyalkylene oxide ethers generally having only a single hydroxyl group (e.g. Klearfac[®], manufacturer: BASF Corp.).

Suitable polyalkylene oxide ethers are, for example, ethers of alkylphenols such as nonylphenol or of branched or unbranched aliphatic alcohols, for example having 6 to 30, preferably having 10 to 20, carbon atoms and in particular of fatty alcohols having 10 to 12 carbon atoms.

The monohydroxylated polyalkylene oxide ethers are generally known or accessible in a manner known per se, especially by alkoxylation of the corresponding alcohols. Preferred alkoxyating agents are ethylene oxide and propylene oxide, which can be reacted with a suitable phosphorus compound individually, in a mixture, in succession or alternately, by means of which alkoxylation products of differing composition, for example having block structures, can result.

The preparation of these phosphoric acid esters is generally known and is carried out, for example, by reaction of the corresponding monofunctional polyalkylene oxide ether with phosphoric acid, diphosphorus pentoxide, polyphosphoric acid or phosphorus oxytrichloride (cf. "Nonionic Surfactants", Martin Schick (Ed.), Marcel Dekker, New York, 1964, Chapter 11, pages 372 - 394).

10

The proportion of the surfactants can be from 0 to 30, preferably from 2 to 15, % by weight.

Suitable antifoams are aliphatic or aromatic monoalcohols having 5 4 to 14, preferably 6 to 10, carbon atoms, such as n-octanol or n-decanol or silicone surfactants.

The proportion of the antifoams in the mixture is normally from 0.5 to 15 and especially from 3 to 8 % by weight.

10

Suitable cosolvents are mineral oils, naturally occurring oils such as rape oil, soybean oil and the methyl esters of carboxylic acids on which these oils are based, such as methyl oleate and rape oil methyl ester, fatty acid esters, especially 15 with C₁-C₄-alkanols and organic solvents such as benzenes or naphthalenes substituted by straight-chain or branched alkyl groups (Shellsol 150®, Shellsol 200® and Solvesso® brands).

The proportion of the cosolvents in the mixture can be from 1 20 to 60 and especially from 5 to 30 % by weight.

Furthermore, the mixture can contain from 0 to 15 and especially from 2 to 10 % by weight of water.

25 Additionally, the mixture can contain one or more carboxylic acids having from 4 to 20, in particular from 6 to 18, carbon atoms such as oleic acid or 2-ethylhexanoic acid and/or one or more of the dicarboxylic acids on which the compounds I are based, e.g. adipic acid, sebacic acid or succinic acid.

30

The proportion of these (di)carboxylic acids in the mixture is from 0 to 30, preferably from 0 to 10, % by weight.

As component d) the plant growth regulating formulations in 35 accordance with the instant invention can additionally contain further known plant growth regulating compounds.

The biological activity of 16,17-dihydro GAs can be improved in a synergistic manner (cf. S.R. Colby, *Weeds* 25: 20-22) by 40 co-applying them with other plant growth regulators. This is particularly observed when using the preferred compounds VI or VII in formulations or spray solutions in combination with other growth retardants, i. e. compounds leading to a reduction of shoot length.

45

Plant growth retardants of practical relevance are, for instance,

- ethephon (standard compound I),
the quaternary ammonium compounds
- chlormequat chloride (standard compound II)
- 5 - mepiquat chloride (standard compound III),
- compounds possessing a N-containing heterocycle
- ancymidol
- flurprimidol
- paclobutrazol
- 10 - uniconazole-P
- inabenfide,
the acylcyclohexanediones
- prohexadione-Ca (standard compound IV)
- trinexapac-ethyl (standard compound V)

15

16,17-dihydro GAs are also suitable to synergize the biological activity of exogenously applied gibberellins, such as gibberellic acid (GA₃ = standard compound VI) or other GAs with immediate biological activity (e.g. GA₄, GA₇). The commercially available

20 GA₃ and GA₄₊₇ have found a large number of applications [for details see Thomson, W.H. (1995) Agricultural Chemicals, Book III, Thomson Publications, Fresno, USA, p. 29-35].

Particularly in graminaceous species, the preferred compounds VI

25 and VII synergize the action of exogenously applied GAs and can, thus, be used to

- increase the yield of malt and decrease the amount of time required for the malting process in the brewing industry
- increase the yield of sugar cane
- 30 - stimulate germination and seedling development in rice, wheat, barley, oats, rye, maize, sorghum, turf grasses and other plant species.

In addition to enhancing the biological performance of 16,17-di-

35 hydro-GAs by using suitable adjuvants for the preparation of formulations, additional optimisation can be achieved by adding further adjuvants to the spray solution (tank mixing). Particularly preferred adjuvants are ammonium sulphate or citric acid added at a concentration of from 0.01 to 0.5 % by weight,

40 preferably 0.05 to 0.2 % by weight.

Examples

Unless otherwise noted, the growth regulating properties of

45 different compounds in different formulations were determined under greenhouse conditions as follows:

12

Test plants were grown in plastic pots of approximately 12.5 cm in diameter in a substrate provided with sufficient nutrients and water. At an appropriate shoot length the plants were treated with aqueous sprays of the formulations. The growth-regulating action observed was confirmed at the end of the experiment by measuring the height of the plants. The results obtained were compared with the growth height of the untreated plants.

In cases in which a reduction in growth height was achieved, a deeper leaf coloration could be observed. The increased chlorophyll content is indicative of an increased rate of photosynthesis, making for bigger yields.

A. Finding of the most preferred compounds

15

Example 1 Shoot length reduction in small grains and rice*

20	Compound	Structure	Shoot length reduction [activity, relative to control] in				
			wheat	barley	oats	rye	rice
	Comp. II	exo-16,17-Dihydro-GA ₅	+ (+)	+	0	++ (+)	/
	Comp. III	endo-16,17-Dihydro-GA ₅	++	++	0	+++	/
25	Comp. IV	Dichloromethano-16,17-dihydro-GA ₅	+	++	+	++	/
	Comp. V	Dichloromethano-16,17-dihydro-GA ₅ -13-acetate	0	0	+	+	/
30	Comp. VI	endo-16,17-Dihydro-GA ₅ -13-acetate	+++	+++	++ (+)	+++	+++
	Comp. VII	exo-16,17-Dihydro-GA ₅ -13-acetate	+++	+++	+++	+++	+++
35	Comp. IX	exo-16,17-Dihydro-GA ₅ -13-n-propyl ether	++ (+)	++ (+)	++	++	++

* 1 % formulations in cyclohexanone : Emulan EL = 4 : 1

40 0 = no effect, + = low activity, ++ = intermediate activity +++ = high activity, / = not tested

In order to obtain results equivalent to compounds VI, VII, and IX in these plant species, dosages approximately 20 times higher have to be used of the standard compounds S1 (ethephon), S4 (prohexadione-Ca) or S5 (trinexapac-ethyl) applied in commercial formulations. In wheat, standard compounds S2 and S3 require

13

dosages of the respective active ingredients approximately 100 times higher. (Standard compounds S2 and S3 are significantly less active in the other plant species.)

5 Example 2 Shoot length reduction in wheat and barley*

10	Compound	Rate [g/ha ai]	Shoot length	
			Wheat [% of con- trol]	Barley [% of control]
15	Comp. IV	1.6	99	99
		3.1	95	100
		6.3	92	98
		12.5	83	87
20	Comp. V	1.6	101	99
		3.1	95	97
		6.3	84	82
		12.5	75	76
25	Comp. VI	1.6	88	95
		3.1	88	84
		6.3	68	65
		12.5	64	53
30	Comp. VII	1.6	90	97
		3.1	73	78
		6.3	70	61
		12.5	64	57

30

*1% formulations in Atlas G 1086 were used.

The results of Examples 1 and 2 demonstrate that compounds VI, VII, and IX are superior to compounds II to V.

35

Examples 3 and 4

Comparison of the plant growth-retarding activity of Compound VII using distinct adjuvants for the preparation of spray solutions
40 or formulations

45

14

Assays with wheat, barley, oat, and rye plants were carried out as described above. The aqueous spray solutions were prepared as follows:

- 5 A Spray solution containing 5 % ethanol (cf. PCT/AU95/00528, Example 39).
 B Spray solution containing 5 % ethanol and 0.1 % Agral 90 (cf. PCT/AU95/ 95/00426, Example 40).
 C Spray solution containing 10 % ethanol and 0.1 % Silwett L
 10 477 (cf. PCT/AU92/00426, Example 11).
 D Spray solution prepared from a 1 % formulation of Compound VII in cyclohexanone : Emulan EL (BASF AG, Ludwigshafen, Germany) = 4:1.
 E Spray solution prepared from a 1 % formulation of Compound
 15 VII in Atlas G 1086 (ICI Surfactants, Fernhurst, Haslemere, Surrey GU27 3JE, GB)
 F Spray solution prepared from a 1 % formulation of Compound VII in ethoxylated castor oil R 15

20 Example 3

25	Preparation of spray solution	Compound VII [g/ha ai]	Shoot length [% of control]	
			Barley	Wheat
30	A	1.5	92	95
		3.0	89	95
		6.0	94	90
		12.0	88	85
35	B	1.5	89	87
		3.0	79	80
		6.0	76	79
		12.0	73	78
40	C	1.5	83	87
		3.0	82	86
		6.0	79	79
		12.0	74	79
45	D	1.5	92	91
		3.0	82	83
		6.0	76	82
		12.0	71	79
	F	1.5	85	85
		3.0	82	83
		6.0	71	79
		12.0	68	75

45

15

As can be seen from Examples 3 and 4, the biological performance of 16,17-dihydro GAS can be particularly enhanced if adjuvants such as Emulan EL, Atlas G 1086 or ethoxylated castor oil R 15 are used for formulation.

5

Example 5

Specific improvement of the biological performance of 16,17-dihydro GAS by ethoxylated castor oil R 15 (R 15)

10	Active Ingredient	Rate [g/ha ai]	Adjuvant	Rate [g/ha ai]	Shoot length [% of control]	
					Barley	Wheat
15	-	-	Ethanol	500	100	100
	-	-	R 15	500	99	98
	Compound VII	2.5	Ethanol	500	99	99
20	Compound VII	2.5	R 15	500	75	73
	Standard Compound S1	250	-	-	95	93
	Standard Compound S1	250	R 15	500	101	92
25	Standard Compound S2	750	-	-	94	83
	Standard Compound S2	750	R 15	500	94	75
30						

The results contained in Example 5 demonstrate that there is a specific and surprising interaction between the 16,17-dihydro GAS and these formulation components since such activity-enhancing effects are not or only at a far lower degree found if equivalent experiments are carried out using standard compounds S1 or S2, respectively.

40 Example 6 Synergistic combinations

To show the synergistic effects of formulations in accordance with the instant invention together with further growth regulating agents, formulations in accordance with the invention were prepared and additional active ingredients added.

16

The results are given in the following tables.

Shoot height reduction in spring barley (cv. "Alexis") 20 days after treatment with combinations of standard compound S2 and 5 compound VII

	Compound	Rate [g/ha a.i.]	% Shoot height reduction		Synergism?
			Actual	Expected	
10	S2	750	11	/	/
	S2	1500	10	/	/
	VII	3	6	/	/
	VII	6	18	/	/
15	S2 + VII	750 + 3	14	16	+
	S2 + VII	750 + 6	40	27	+
	S2 + VII	1500 + 3	20	15	+
	S2 + VII	1500 + 6	40	26	+

20 Shoot height reduction in spring wheat (cv. "Ralle") 21 days after treatment with combinations of standard compound S4 and compound VII

	Compound	Rate [g/ha a.i.]	% Shoot height reduction		Synergism?
			Actual	Expected	
25	S4	30	6	/	/
	S4	60	15	/	/
	VII	3	8	/	/
	VII	6	21	/	/
30	S4 + VII	30 + 3	16	14	(+)
	S4 + VII	30 + 6	33	26	+
	S4 + VII	60 + 3	42	22	+
	S4 + VII	60 + 6	49	33	+

35

40

45

17

Shoot height reduction in spring barley (cv. "Alexis") 20 days after treatment with combinations of standard compound S4 and compound VII

5	Compound	Rate [g/ha a.i.]	% Shoot height reduction		Synergism?
			Actual	Expected	
	IV	30	10	/	/
	IV	60	27	/	/
	6	3	6	/	/
10	6	6	18	/	/
	S4 + VII	30 + 3	29	15	+
	S4 + VII	30 + 6	45	26	+
	S4 + VII	60 + 3	39	31	+
15	S4 + VII	60 + 6	47.5	40	+

Shoot height reduction in spring wheat (cv. "Ralle") 13 days after treatment with combinations of standard compound S5 and compound VII

20	Compound	Rate [g/ha a.i.]	% Shoot height reduction		Synergism?
			Actual	Expected	
	S5	30	14	/	/
	VII	6	18	/	/
25	S5 + VII	30 + 6	32	17	+

16,17-dihydro GAS are also suitable to synergize the biological activity of exogenously applied gibberellins, such as gibberellic acid (GA₃ = standard compound S6) or other GAS with immediate biological activity (e.g. GA₄, GA₇). The commercially available GA₃ and GA₄₊₇ have found a large number of applications [for details see Thomson, W.H. (1995) Agricultural Chemicals, Book III, Thomson Publications, Fresno, USA, p. 29-35].

35

Particularly in graminaceous species, the preferred compounds VI and VII synergize the action of exogenous GAS and can, thus, be used to

40 - increase the yield of malt and decrease the amount of time required for the malting process in the brewing industry

- increase the yield of sugar cane

45 - stimulate germination and seedling development in rice, wheat, barley, oats, rye, maize, sorghum, turf grasses and other plant species.

The synergizing effect of compound VII on shoot elongation in rice is shown in the results of Example 7. The shoot growth-promoting effect of GA₃ could clearly be raised by compound VII which, applied alone, reduces shoot elongation.

5

Example 7

Enhanced shoot growth of rice seedlings [cvs. "Koshihikari" (tall), "Tan-ginbozu" (dwarf), "Waito C" (dwarf)] 7 days after treatment with combinations of compound 6 with GA₃ ^d

10

Compound(s)	Rate [g/ha a.i.]	Koshihikari [% of Con- trol] ^a	Tan-ginbozu [% of Con- trol] ^b	Waito C [% of Con- trol] ^c
VII	12.5	75	74	80
VII	25.0	75	74	80
VII	50.0	75	74	80
S6	1.25	109	128	115
S6	2.50	112	160	143
S6	5.00	124	186	190
VII S6	12.5 1.25	118	170	167
VII S6	12.5 2.50	132	191	195
VII S6	12.5 5.00	147	229	230
VII S6	25.0 1.25	129	197	184
VII S6	25.0 2.50	141	223	207
VII S6	25.0 5.00	149	229	230
VII S6	50.0 1.25	129	202	184
VII S6	50.0 2.50	135	213	218
VII S6	50.0 5.00	155	234	253

a 100 % = 17.4 cm / shoot length at foliar treatment: 12.5 cm

b 100 % = 9.4 cm / shoot length at foliar treatment: 6.0 cm

c 100 % = 8.7 cm / shoot length at foliar treatment: 5.0 cm

d 1 kg/ha of ammonium sulphate added to the spray solution

Claims:

1. Plant growth regulating preparations comprising
 - 5 a) 0.1 to 20 % by weight of a 16,17-dihydro gibberellin of general formulae Ia or Ib
 - b) up to 99.9 % by weight of an formulation additive selected from the group consisting of
 - 10 b1) the reaction products of triglycerides based on carboxylic acids having 2 to 30 carbon atoms and ethylene oxide and/or propylene oxide in the presence of a base, and/or
 - 15 b2) fatty acid esters of sugar alcohol polyethoxylates
 - c) up to 50 % by weight of an organic solvent
 - 20 d) 0.1 to 50 % by weight of a formulation auxiliary different from b1 and b2,
 - e) up to 50 % by weight of additional plant growth regulating compounds.
- 25 2. Plant growth regulating formulations as defined in claim 1 comprising as component 1 exo-16,17-dihydro-GA₅-13-acetate, endo-16,17-dihydro-GA₅-13-acetate, exo-16,17-dihydro-GA₅-13-C₁-C₄-alkyl ether or endo-16,17-dihydro-GA₅-13-C₁-C₄-alkyl ether or mixtures thereof.
 - 30
3. Plant growth regulating formulations as defined in claim 1 or 2, comprising as component b a compound b1.
- 35 4. Plant growth regulating formulations as defined in any of claims 1 or 2 comprising as component b a compound b2.
5. Plant growth regulating formulations as defined in any of the preceding claims, comprising as component e) a compound
 - 40 selected from the group consisting of chlormequat chloride, mepiquat chloride, prohexadione-Ca, trinexapac ethyl and ethephon or mixtures thereof.
6. Plant growth regulating formulations comprising GA₃ or a
 - 45 mixture of GA₄ and GA₇ as component e).

20

7. Use of the plant growth regulating formulations as defined in any of claims 1 to 5 for the retardation of plant growth.
8. Use of the plant growth regulating formulations as defined in claim 6 to stimulate shoot growth in graminaceous plant species.

10

15

20

25

30

35

40

45

INTERNATIONAL SEARCH REPORT

Inter .nal Application No
PCT/EP 99/04592

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N45/00 A01N25/30 //(A01N45/00, 57:20, 45:00, 43:40, 37:42, 33:12)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 93 03616 A (EVANS LLOYD THOMAS ;WHITFIELD KING RODERICK (AU); MANDER LEWIS NOR) 4 March 1993 (1993-03-04) cited in the application page 1 -page 9, paragraph 2 page 11, paragraph 3 page 13, last paragraph -page 14, paragraph 1 page 18, last paragraph -page 19, paragraph 1; example 11 ---	1-8
Y	GB 862 849 A (MERCK & CO., INC.) page 1 -page 3, line 8; claims --- -/--	1-8

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

22 October 1999

Date of mailing of the international search report

03/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Muellners, W

INTERNATIONAL SEARCH REPORT

Inter inal Application No
PCT/EP 99/04592

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 06090 A (UNIV AUSTRALIAN ;COMMW SCIENT IND RES ORG (AU); PHARIS RICHARD PER) 29 February 1996 (1996-02-29) cited in the application page 1 -page 6 page 11, line 22 - line 27; examples 39,40 ---	1-8
A	GB 930 078 A (R.J.GEARY) page 2, line 16 - line 95 page 5, line 62 - line 100 ---	1-8
A	US 5 163 993 A (SHAFFER WARREN E ET AL) 17 November 1992 (1992-11-17) column 1, line 58 -column 2, line 9 column 3, line 10 -column 4, line 17 ---	1-8
A	EP 0 232 755 A (MITSUBISHI GAS CHEMICAL CO) 19 August 1987 (1987-08-19) page 2, line 10 -page 3, line 27 page 8, line 1 - line 35 formulation examples 1 and 2 -----	1-8

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 99/04592

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9303616 A	04-03-1993	AU 665961 B AU 2444092 A DE 69228188 D DE 69228188 T EP 0599939 A ES 2129451 T NZ 243973 A US 5532206 A	25-01-1996 16-03-1993 25-02-1999 22-07-1999 08-06-1994 16-06-1999 27-09-1994 02-07-1996
GB 862849 A		NONE	
WO 9606090 A	29-02-1996	AU 703725 B AU 3248395 A CA 2198355 A EP 0777662 A JP 10504549 T US 5767042 A	01-04-1999 14-03-1996 29-02-1996 11-06-1997 06-05-1998 16-06-1998
GB 930078 A		BE 584219 A FR 1245061 A NL 121956 C NL 245018 A US 3038794 A	27-01-1961 12-06-1962
US 5163993 A	17-11-1992	NONE	
EP 0232755 A	19-08-1987	JP 1863243 C JP 5078522 B JP 62167704 A US 4799950 A	08-08-1994 29-10-1993 24-07-1987 24-01-1989

